Ground-based capabilities for general astrophysics and exoplanets science

Olivier Guyon

University of Arizona
Subaru Telescope, National Astronomical Observatory of Japan, National Institutes for Natural Sciences
Astrobiology Center, National Institutes for Natural Sciences
JAXA

HabEx meeting, Aug 3, 2016

HabEx Uniqueness

HabEx unique capabilities (not accessible from ground):

- Largest UV-Opt astronomical aperture in space
- Spectral coverage: UV not accessible from ground, continuous NIR coverage
- Angular resolution in UV .. and optical ?
- Ultra-high contrast
- High stability
  - → astrometry
  - → precision photometry, spectroscopy

Major ground facilities (Opt-NIR)

ELTs coming online in mid-2020s:

- E-ELT: 39m aperture, Chile
- TMT: 30m, Hawaii(?)
- GMT: 25m, Chile

8-m class survey telescope/instruments:

- Imaging: LSST
- Spectroscopy: Subaru-PFS

+ other dedicated survey facilities (LAMOST, PAN-STARRS PTF etc…)

Wide FOV optical imaging: Subaru HSC











8m aperture, 1.5deg diam FOV
104 4kx2k CCDs

LSST





8m aperture, 3.5deg diam FOV
189 4kx4k CCDs

Subaru Prime Focus Spectrograph



2,400 fibers over 1.3 deg diam FOV
0.38 – 1.26 um

Increased image quality over moderate FOV in near-IR with GLAO

example: ULTIMATE-Subaru

ULTIMATE key specifications: Ground-Layer AO

Field of View	15 arcmin
Focus	Cassegrain
Corrector	Adaptive secondary, 33 actuators across the diameter
Sky Coverage	Almost all sky

FWHMs at 'Moderate' condition

	GLAO	Natural
K-band	0.20"	0.44"
H-band	0.23"	0.49"
J-band	0.27"	0.51"
R-band	0.41"	0.65"

ULTIMATE-Subaru Instruments



ELTs







E-ELT first light instruments



E-ELT – First Light instruments

MAORY + MICADO
(Multi-conjugate Adaptive Optics RelaY for the E-ELT)
(Multi-AO Imaging Camera for Deep Observations)
0.8 – 2.4 um
diffraction-limited imaging (6 – 12 mas)
R=8000 spectroscopy

HARMONI
(High Angular Resolution Monolithic Optical and Near-infrared Integral field spectrograph)
0.47 – 2.45 um
IFS
R = 3000 – 20,000 spectroscopy

METIS
10um imaging and spectroscopy
high contrast imaging

E-ELT instruments

Exoplanet science:
CODEX (precision RV), METIS (10um imaging) and EPICS/PCS (direct imaging, NIR)





Multi-Conjugate Adaptive Optics (MCAO)

Concept: Use several DMs conjugated at different altitudes to perform correction over a wide fielf of view



Gemini South MCAO system



EAGLE
(ESO E-ELT project)

MCAO system with 20 fields
0.8-2.45 um



1 Shutter
2 Laser Guide Star Sensing System
3 Pick-off System (Focal Plane)
4 Target Reimaging and Magnification System (including Deformable Mirror)
5 Integral Field Unit and Spectrograph System



TMT instruments

Wide FOV imager
NIR, 30”

Near-IR imaging (0.83-2.4um)
+ IFU, R=4000-10000

High resolution spectro
R=50,0000
0.31-1.1 um
(Seeing limited)

AO system
+ IRMS multi-slit spectro

high contrast
imaging



Visible light MOS
40 sq ' FOV
R=1000-8000
0.31-1.1 um
(Seeing limited)

High res spectro
mid-IR
R=100,000
4.5” slit length
10” imaging FOV

Multiple IFUs
0.8-2.5 um
3” FOV per IFU
MOAO-fed

AO-fed TMT instruments (First light)



IRIS



IRMS



GMT instruments: Optical spectroscopy





0.35 – 0.95 um
R: 19,000 – 108,000

Exoplanets: precision radial velocity, transit spectroscopy
Stellar physics
Galaxies, cosmology

GMACS

0.35 – 0.95 um
R = 1,000 – 6,000

Multi-objects
(100s of spectra)

Galaxies, cosmology



GMT instruments: NIR spectroscopy

Near-IR IFU and Adaptive Optics Imager – GMTIFS



0.9 – 2.5um IFU
R = 5,000 to 10,000
FOV: 20”x20” (imager), 0.3”x0.6”, 2.2”x4.5” (IFU)
5mas – 50mas resolution

IR Echelle Spectrograph – GMTNIRS



1.1 – 5.3um
R = 50,000 to 75,000

Exoplanet capabilities - RV

Optical RV is making steady progress and will reveal lower mass planets

Mix of high precision instruments on largest telescopes (for example CODEX on E-ELT), and dedicated robotic RV facilities

Near-IR RV will identify habitable planets around nearby M-type stars. First instruments being deployed **NOW**



SPIROU @ CFHT



Habitable Zones within 5 pc (16 ly): Astrometry and RV Signal Amplitudes for Earth Analogs



Exoplanet capabilities – Direct imaging

10um imaging: METIS @ E-ELT
→ thermal emission from Earth-size planets & larger around nearby stars

Near-IR / visible Extreme-AO:

8m telescopes: spectroscopy of self-luminous planets (young Jupiters)
.. and possibly reflected light imaging of nearby large planets

ELTs: detailed characterization of giant planets
Reflected light imaging and spectroscopy of Earths around M (& K?) type stars

Spectroscopic characterization of Earth-sized planets with ELTs




Habitable Zones within 5 pc (16 ly)
Star Temperature [K]
1 Gliese 1245 A
2 Gliese 1245 B
3 Gliese 674
4 Gliese 440 (white dwarf)
5 Gliese 876 (massive planets in/near HZ)
6 Gliese 1002
7 Gliese 3618
8 Gliese 412 A
9 Gliese 412 B
10 AD Leonis
11 Gliese 832
LP944-020
SRC 1845-6357 A
DEN1048-3956
Wolf 424 A
Van Maanen's Star
Ross 614 B
LHS 292
DX Cnc
Teegarden's star
TZ Arietis
CN Leonis
YZ Ceti
Wolf 424 B
UV Ceti
GJ 1061
LHS380
BL Ceti
EZ Aqu A,B & C
Proxima Cen
Ross 248
Kruger 60 B
Gl15 B
Barnard's Star
Ross 154
Ross 128
Ross 614 A
Kapteyn's star
40 Eri C
Procyon B
Gl725 B
Wolf 1061
40 Eri B
Gl725 A
Gliese 1
Lalande 21185
Gl15 A
Sirius B
Lacaille 9352
Kruger 60 A
Luyten's star
Gliese 687
AX Mic
61 Cyg A
Groombridge 1618
61 Cyg B
Eps Indi
Eps Eri
Tau Ceti
40 Eri A
α Cen B
α Cen A
Procyon A
Sirius
A
F
G
K
M
log Contrast
distance (pc)
Circle diameter indicates angular size of habitable zone
Circle color indicates stellar temperature (see scale right of figure)
Contrast is given for an Earth analog receiving the same stellar flux as Earth receives from Sun (reflected light)


Habitable Zones within 5 pc (16 ly)
Star Temperature [K]
1 Gliese 1245 A
2 Gliese 1245 B
3 Gliese 674
4 Gliese 440 (white dwarf)
5 Gliese 876 (massive planets in/near HZ)
6 Gliese 1002
7 Gliese 3618
8 Gliese 412 A
9 Gliese 412 B
10 AD Leonis
11 Gliese 832
Radial Velocity (near-IR)
ELT imaging (near-IR)
Space astrometry
ELT imaging (thermal-IR)
LP944-020
SRC 1845-6357 A
DEN1048-3956
Wolf 424 A
Van Maanen's Star
DX Cnc
Ross 614 B
Teegarden's star
LHS 292
CN Leonis
YZ Ceti
UV Ceti
BL Ceti
GJ 1061
Wolf 424 B
LHS380
Proxima Cen
EZ Aqu A,B & C
Ross 248
Ross 154
Ross 128
Gl15 B
Kruger 60 B
Barnard's Star
Ross 614 A
Kapteyn's star
40 Eri C
40 Eri B
Procyon B
Gl725 B
Wolf 1061
Gl725 A
Gliese 1
Lalande 21185
Gl15 A
Sirius B
Lacaille 9352
Kruger 60 A
Luyten's star
Gliese 687
AX Mic
61 Cyg A
61 Cyg B
Groombridge 1618
Eps Indi
Eps Eri
Tau Ceti
40 Eri A
α Cen B
α Cen A
Procyon A
Sirius
A
F
G
K
M
9000
8000
7000
6000
5000
4000
3000
log Contrast
-5
-6
-7
-8
-9
-10
-11
-12
0
1
2
3
4
5
distance (pc)
Circle diameter indicates angular size of habitable zone
Circle color indicates stellar temperature (see scale right of figure)
Contrast is given for an Earth analog receiving the same stellar flux as Earth receives from Sun (reflected light)

Summary

Ground-based telescopes will provide excellent visible light wide field imaging and MOS in visible and near-IR
Low resolution spectroscopy (IFS) and high resolution (R~100,000)

Adaptive optics will provide diffraction limited imaging anywhere in the sky at near-IR, and around bright sources (m~12 and brighter) in visible light

AO may provide nearly full sky coverage diffraction limited images in visible light (see Keck visible AO development plans)

Exoplanets:

RV (especially nearIR) will identify habitable planets

Spectroscopic characterization with direct imaging: Giant planets with 8m telescopes, Earth-like planets with ELTs

Nearest M (&K)-type stars provide opportunities for direct imaging + RV/astrometry